

## Providing a "quick wash" to oil contaminated wildlife using magnetic particle technology (MPT)

**John D. Orbell, Stephen W. Bigger, Lawrence N. Ngeh, Hien Van Dao and Kasup Munaweera**

School of Engineering & Science, Victoria University, PO Box 14428, Melbourne 8001, AUSTRALIA

**Peter Dann, Rosalind Jessop and Margaret Healy**

Research Department, Phillip Island Nature Park, Victoria 3922, Australia

HEALTH,  
ENGINEERING  
AND SCIENCE

WWW.VU.EDU/AU/HES


VICTORIA  
UNIVERSITY

A NEW  
SCHOOL OF  
THOUGHT

## How can MPT be applied to environmental remediation and wildlife rehabilitation?

But the science is very challenging.....

A very simple concept.....



Development of the particles themselves – different materials, surface variation, size distributions, coatings etc....

Optimizing the particles with respect to different oils and substrates, such as rock surfaces, feathers, plumage, fur, live animals....

Development of the 'magnetic harvester' – field strength, on/off mechanism, design features etc.....

Development of methodologies and protocols for testing ....

Moving proof of principle 'lab work' to the field....

Waste disposal for different scenarios.....

Logistics and costs....

Potential spin-offs.....


## Particle development .....

Various particles continue to be explored and tested including -


Iron particles coated with a hydrophobic oil-adsorbing material such as PVC or PE.

Different grades of finely divided iron powder with different surface characteristics.


Iron particles with "nanoscale surface roughness" resulting in "super-hydrophobic" properties when coated with materials such as polydimethylsiloxane (PDMS).



PVC coated iron pellet




Different grades of iron powder



AFM imaging of iron particle surface

## Optimizing the particles themselves.....


Comparative removal of oil from feathers for eight different grades of iron powder. MH300.29 ("spongy superfine") achieves 100% within experimental error.



H. V. Dao, L.N. Ngeh, S.W. Bigger and J.D. Orbell, 2005, "The achievement of 100% removal of oil from feathers employing magnetic particle technology", *J. Environ. Eng.*, Vol. 132, No. 5, 555-559.

## Different oil types and ambient conditions.....


Removal of tarry/weathered contamination from feathers -



J.D. Orbell, H.V. Dao, L.N. Ngeh, S.W. Bigger, M. Healy, R. Jessop and P. Dann, 2005, "Acute temperature dependency in the cleansing of tarry feathers utilizing magnetic particles", *Environ. Chemistry Letters*, 3(1), 25-27.


## Why does this happen?

The transfer of oil from the surface of the tarry residue onto the iron particles represents competitive adsorption between two different surfaces:



The experimental design allows the equilibrium constant for the process to be estimated at different temperatures and at different stages of removal.

This allows the Van't Hoff Equation,  $\ln K = -\Delta H^\circ/RT + \Delta S^\circ/R$ , to be applied.



Overall we can say that:

The process is *endothermic*. This favours transfer of oil onto the iron as the temperature increases.

The process is *highly entropy driven*.

J.D. Orbell, H.V. Dao, L.A. Maher, L.N. Ngeh, S.W. Bigger, M. Healy, R. Jessop and P. Dann, 2006, "Removal of petroleum tar from bird feathers utilizing magnetic particles", *Environ. Chemistry Letters*, 4, 111-113.

## Optimizing the magnetic harvester....

"Horses for courses".....



The 'magnetic tester' – suitable for laboratory testing and proof of principle studies. Requires two hands to operate.



Trial electromagnetic device – insufficient magnetic field strength.




One-handed device powered by compressed air – too cumbersome for field work




The "magnetic wand" designed for a "quick clean (wash)". Strong localized magnetic field – non magnetic tip.

## Different substrates.....


From feather clusters



From rock surface



From Mammalian fur



From plumage – whole bird models



Clean Little Penguin carcass



50% coverage of bunker oil



After 5 treatments




After 14 treatments

J.D. Orbell, H.V. Dao, J. Kapadia, L.N. Ngh, S.W. Bigger, M. Healy, R. Jessop and P. Dann, 2007, "An Investigation into the removal of oil from rock utilizing magnetic particle technology", *Marine Pollution Bulletin*, 54, 1958 – 1961.


J.D. Orbell, L.N. Ngh, S. W. Bigger, M. Zabinskas, M. Zheng, M. Healy, R. Jessop and P. Dann, 2004, "Whole-Bird Models for the Magnetic Cleansing of Oiled Feathers", *Marine Pollution Bull.*, 48, 336-340.

## Development of specific methodologies.....

For example, determining the % coverage by mass for controlled experiments.




For example: for a bird with a body weight of 620 g the weight of oil required for saturation (100 % coverage) is  $(0.2339 \times 620) - 8.6546 = 136.2$  g. Therefore, 50% coverage by weight requires  $136.4 \times 0.5 = 68.2$  g of oil to be applied.



Diesel required for 50% coverage


## Systematic feasibility experiments - the potential for a "quick wash"



100% diesel coverage of Little Penguin plumage – a "worst case" scenario

J.D. Orbell, H.V. Dao, L.N. Ngh and S.W. Bigger, (2008), "An investigation into the feasibility of applying magnetic particle technology to the cleansing of oiled wildlife in the field". Report prepared for the Australian Maritime Safety Authority (National Plan Environment Working Group) and the Phillip Island Nature Park.

## % coverage experiments (diesel)



Magnetic tester; iron powder grade MH300.29; no pre-conditioner

J.D. Orbell, H.V. Dao, L.N. Ngh, S.W. Bigger, M. Healy, R. Jessop and P. Dann, 2006, "Magnetic cleansing of weathered/tarry oiled feathers - the role of pre-conditioners", *Marine Pollution Bulletin*, 52, 1591-1594.

## "In the field" scenario – some conservative numbers

Based on 100 birds for 1 (blue) treatment and 2 (red) treatments



Number of birds = 100 (100)

Average coverage = 50% (50%)

Treatments per bird = 1 (2)

Average time for treatment(s) = 2.5 min (5 min)

Average mass of powder per bird = 35 g (68 g)

Average % oil removal = 25% (40%)

Average mass of oil removed per bird = 17 g (27 g)

Number of 2-person teams = 2 (2)

Total person hours required = 4.2 hrs (8.4 hrs)

Total time per 2-person team = 2.1 hrs (4.2 hrs)

Number of birds per 2-person team = 50 (50)

Total mass of powder = 3.5 kg (6.8 kg)

Total mass of oil removed = 1.7 kg (2.7 kg)

Total mass of waste = 5.2 kg (9.5 kg)

Mass of powder per 2-person team = 1.75 kg (3.4 kg)

Mass of oil per 2-person team = 0.85 kg (1.35 kg)

Mass of waste per 2-person team = 2.6 kg (4.75 kg)

### "Holding bay" scenario – some conservative numbers

Based on 1000 birds for 1 (blue) treatment and 2 (red) treatments



Number of birds = 1000 (1000)  
 Average coverage = 50% (50%)  
 Treatments per bird = 1 (2)  
 Average time per treatment(s) = 2.5 min (5 min)  
 Average mass of powder per bird = 35 g (68 g)  
 Average % oil removal = 25% (40%)  
 Average mass of oil removed per bird = 17 g (27 g)  
 Number of 2-person teams = 10 (10)



Total person hours required = 42 hrs (84 hrs)  
 Total time per 2-person team = 4.2 hrs (8.4 hrs)  
 Number of birds per 2-person team = 100 (100)  
 Total mass of powder = 35 kg (68 kg)  
 Total mass of oil removed = 17 kg (27 kg)  
 Total mass of waste = 52 kg (95 kg)  
 Mass of powder per 2-person team = 3.5 kg (6.8 kg)  
 Mass of oil per 2-person team = 1.7 kg (2.7 kg)  
 Mass of waste per 2-person team = 5.2 kg (9.5 kg)


### Some estimated costs for both scenarios

100 Birds/2 Teams <sup>3</sup>	One treatment	Two treatments
Iron powder at \$5.50/kg	3.5 kg x 5.5 = \$19	6.8 kg x 5.5 = \$37
Waste disposal at \$8/kg	5.2 kg x 8 = \$42	9.5 kg x 8 = \$76
Personnel at \$25 per hour	2.1 hrs x 4 x 25 = \$210	4.2 hrs x 4 x 25 = \$420
<b>Total</b>	<b>\$271</b>	<b>\$533</b>
<b>Cost per bird</b>	<b>\$2.71</b>	<b>\$5.33</b>


1000 Birds/10 Teams	One treatment	Two treatments
Iron powder at \$5.50/kg	35 kg x 5.5 = \$193	68 kg x 5.5 = \$374
Waste disposal at \$8/kg	52 kg x 8 = \$416	95 kg x 8 = \$770
Personnel at \$25 per hour	4.2 hrs x 20 x 25 = \$2100	8.4 hrs x 20 x 25 = \$4,200
<b>Total</b>	<b>\$2,709</b>	<b>\$5,344</b>
<b>Cost per bird</b>	<b>\$2.71</b>	<b>\$5.34</b>

Latest estimate of equipment with "wand" = \$600/team team

### Developing equipment, techniques and logistical data for a quick wash – utilizing the "wand".....



Non-magnetic tip allows oil laden particles to be readily disposed of.



### The Animal Rehabilitation Technology (ART) Group



Phillip Island Research Centre  
 Peter Dann, Roz Jessup, Marg Healy



#### Victoria University

John Orbell, Stephen Bigger, Lawrence Ngeh, Leroy Godhino, Sally Ryan, Hien Van Dao, Ali Abdrahman, David Thompson, Lauren Maher, Eee Kai Tan, Manfred Zabinkas, Meris Zheng, Leroy Godhino, Jignesh Kapadia, Kasup Munawerra



Professor Orbell (front left) and colleagues Ali Abdrahman, Sally Ryan, Hien Van Dao, Dr. Lawrence Ngeh and Associate Professor Manfred Bigger (front right).